



CONCUSSION

AWARENESS & MANAGEMENT



THINGS YOU NEED TO KNOW

CONCUSSION: Signs & Symptoms

- Confusion and Disorientation
- Loss of Consciousness
- Headache
- Dizziness
- Nausea and Vomiting
- Loss of Balance
- Double Vision or Fuzzy Vision
- Ringing in the Ears
- Slow or Slurred Speech
- Seeing "Stars"
- Feeling Stunned or Dazed
- Emotional or Personality Changes

CONCUSSION: Management & Rehabilitation

An athlete should never return to play while symptomatic.

When a player shows ANY SYMPTOMS or SIGNS of a concussion...

✗ The player should not be allowed to return to play in the current game or practice.

✗ The player should not be left alone, regular monitoring for deterioration is essential.

✓ The player should be medically evaluated.

CONCUSSION: Guidelines for Coaches, Players, Parents & Officials

- Concussion is a Brain Injury
- You do not have to lose consciousness to have a concussion
- Symptoms are often subtle
- Wear properly fitted protective equipment
- **The head (helmet/facemask) should never be used to make initial contact with another player**
- A concussion may be caused by a direct blow to the head, face, neck, or anywhere else that causes a severe and sudden movement to the head/neck
- Medical Clearance by an appropriate physician is mandatory before return to play

CONCUSSION: Follow these 6 steps before returning to play.

Players must be asymptomatic for 24 hours BETWEEN each step.

- 1.** No activity, complete rest. Once asymptomatic, proceed to step two
- 2.** Light aerobic exercise such as walking or stationary cycling
- 3.** Sport-specific training. (e.g. running in football, skating in hockey)
- 4.** Non-contact training drills. May start progressive resistance training
- 5.** Full-contact training after medical clearance
- 6.** Game Play



thinkfirst

